

ON WEIGHTED POINCARÉ INEQUALITIES

BARTŁOMIEJ DYDA AND MORITZ KASSMANN

ABSTRACT. The aim of this note is to show that Poincaré inequalities imply corresponding weighted versions in a quite general setting. The proof is short and does not involve covering arguments.

1. INTRODUCTION

Let (X, ρ) be a metric space with a positive σ -finite Borel measure dx , we will write $|E| = \int_E dx$ for the measure of a Borel set $E \subset X$. We fix some point $x_0 \in X$ and set $B_r = \{x \in X : \rho(x, x_0) < r\}$, $\overline{B}_r = \{x \in X : \rho(x, x_0) \leq r\}$.

We call a function $\phi : B_1 \rightarrow [0, \infty)$ an *admissible weight*, if ϕ is a radial function, i.e. $\phi = \Phi(\rho(\cdot, x_0))$ and its profile Φ is nonincreasing and right-continuous with left-limits. We assume that ϕ is not identically zero on $B_1 \setminus \overline{B}_{1/2}$.

For any admissible weight ϕ there exists a positive, non-zero σ -finite Borel measure ν on $(\frac{1}{2}, 1]$, such that

$$(1) \quad \phi(x) = \int_{\rho(x, x_0) \vee 1/2}^1 \nu(dt) = \int_{1/2}^1 \chi_{B_t}(x) \nu(dt), \quad x \in B_1 \setminus \overline{B}_{1/2}.$$

(Note that we put $\int_a^b f(t) \nu(dt) := \int_{(a, b]} f(t) \nu(dt)$.)

For a function u we denote by

$$u_E = \frac{1}{|E|} \int_E u(x) dx$$

the mean of u over the set E , and by

$$u_E^\phi = \frac{\int_E u(x) \phi(x) dx}{\int_E \phi(x) dx}$$

the mean of u over the set $E \subset B_1$ with respect to the weight function ϕ .

Our main result is the following:

Date: October 2, 2012.

2010 Mathematics Subject Classification. Primary 35A23; Secondary 26D10, 26D15.

Key words and phrases. Dirichlet forms, Sobolev spaces, Poincaré inequality.

Both authors have been supported by the German Science Foundation DFG through SFB 701. The first author was additionally supported by MNiSW grant N N201 397137.

Theorem 1. *Let $1 \leq p < \infty$ and let ϕ be an admissible weight with $\phi = \Phi(\rho(\cdot, x_0))$. Let $F : L^p(X) \times (\frac{1}{2}, 1] \rightarrow [0, \infty]$ be a functional satisfying*

$$(2) \quad F(u + a, r) = F(u, r), \quad a \in \mathbb{R},$$

$$(3) \quad \int_{B_r} |u(x) - u_{B_r}|^p dx \leq F(u, r),$$

for every $r \in (\frac{1}{2}, 1]$ and every $u \in L^p(X)$. Then for $M = \frac{8^p |B_1|}{|B_{1/2}|} \frac{\Phi(0)}{\Phi(1/2)}$

$$(4) \quad \int_{B_1} |u(x) - u_{B_1}^\phi|^p \phi(x) dx \leq M \int_{1/2}^1 F(u, t) \nu(dt)$$

for every $u \in L^p(B_1)$, where ν is as in (1).

By choosing the functional F appropriately, (4) becomes a Poincaré inequality with weight ϕ , see Section 3. Such inequalities have been studied extensively because of their importance for the regularity theory of partial differential equations, see the exposition in [5].

2. PROOF

Lemma 2. *Let Ω be a finite measure space and $p \geq 1$. Assume $f \in L^p(\Omega)$ with $\int_\Omega f = 0$. Then*

$$\|f + a\|_{L^p(\Omega)} \geq \frac{1}{2} \|f\|_{L^p(\Omega)}$$

for every $a \in \mathbb{R}$.

Proof. We may assume $a > 0$. Then

$$\int_{\Omega \cap \{f > 0\}} |f + a|^p \geq \int_{\Omega \cap \{f > 0\}} |f|^p \quad \text{and} \quad \int_{\Omega \cap \{f < -2a\}} |f + a|^p \geq 2^{-p} \int_{\Omega \cap \{f < -2a\}} |f|^p.$$

Furthermore, since $\int_{\Omega \cap \{f \leq 0\}} |f| = \int_{\Omega \cap \{f > 0\}} |f|$, we obtain

$$\int_{\Omega \cap \{-2a \leq f \leq 0\}} |f|^p \leq (2a)^{p-1} \int_{\Omega \cap \{-2a \leq f \leq 0\}} |f| \leq (2a)^{p-1} \int_{\Omega \cap \{f > 0\}} |f| \leq 2^{p-1} \int_{\Omega \cap \{f > 0\}} |f + a|^p,$$

where we use $a^{p-1}b \leq (b+a)^{p-1}(b+a)$ for positive a, b . Combining these observations we obtain the result. \square

Proof of Theorem 1. First we observe that it is enough to prove that

$$(5) \quad \int_{B_1} |u(x) - u_{B_1}^{\tilde{\phi}}|^p \tilde{\phi}(x) dx \leq \frac{2^{2p} |B_1|}{|B_{1/2}|} \int_{1/2}^1 F(u, t) \nu(dt),$$

where $\tilde{\phi}(x) = \phi(x) \wedge \Phi(\frac{1}{2})$. Indeed, we have

$$\frac{\Phi(\frac{1}{2})}{\Phi(0)} \phi(x) \leq \phi(x) \wedge \Phi(\frac{1}{2}) \leq \phi(x).$$

Hence if (5) holds, then

$$\begin{aligned} \int_{B_1} |u(x) - u_{B_1}^{\tilde{\phi}}|^p \tilde{\phi}(x) dx &\geq \frac{\Phi(\frac{1}{2})}{\Phi(0)} \int_{B_1} |u(x) - u_{B_1}^{\tilde{\phi}}|^p \phi(x) dx \\ &\geq \frac{\Phi(\frac{1}{2})}{\Phi(0)} 2^{-p} \int_{B_1} |u(x) - u_{B_1}^{\phi}|^p \phi(x) dx, \end{aligned}$$

where in the last line we have used Lemma 2.

Now we prove (5). To simplify the notation, we assume that $\phi(x) = \Phi(\frac{1}{2})$ for $x \in B_{1/2}$, so that $\tilde{\phi} = \phi$.

Because of (2), by subtracting a constant from u , we may and do assume that $u_{B_1}^{\phi} = 0$, which means that

$$(6) \quad 0 = \int_{B_1} u(x) \phi(x) dx = \int_{1/2}^1 \int_{B_t} u(x) dx \nu(dt) = \int_{1/2}^1 u_{B_t} |B_t| \nu(dt).$$

We start from the integral on the right hand side of (4) and use (3)

$$\begin{aligned} R &:= \int_{1/2}^1 F(u, t) \nu(dt) \geq \int_{1/2}^1 \int_{B_t} |u(x) - u_{B_t}|^p dx \nu(dt) \\ &= \frac{1}{2} \int_{1/2}^1 \int_{B_t} |u(x) - u_{B_t}|^p dx \nu(dt) + \frac{1}{2} \int_{B_1} \int_{1/2}^1 |u(x) - u_{B_t}|^p \chi_{B_t}(x) \nu(dt) dx \\ &=: I_1 + I_2 \end{aligned}$$

(In fact $I_1 = I_2$, but we treat them differently.) We now deal with the inner integral in I_2 . For $x \in B_{1/2}$ we have

$$\int_{1/2}^1 |u(x) - u_{B_t}|^p \chi_{B_t}(x) \nu(dt) \geq \frac{1}{|B_1|} \int_{1/2}^1 |u(x) - u_{B_t}|^p |B_t| \nu(dt).$$

Since $\int_{1/2}^1 u_{B_t} |B_t| \nu(dt) = 0$, by Lemma 2 we obtain

$$\int_{1/2}^1 |u(x) - u_{B_t}|^p |B_t| \nu(dt) \geq 2^{-p} \int_{1/2}^1 |u_{B_t}|^p |B_t| \nu(dt).$$

Therefore

$$I_2 \geq \frac{2^{-p}}{2|B_1|} \int_{B_{1/2}} \int_{1/2}^1 |u_{B_t}|^p |B_t| \nu(dt) dx = \frac{2^{-p}|B_{1/2}|}{2|B_1|} \int_{1/2}^1 |u_{B_t}|^p |B_t| \nu(dt).$$

Using inequality $|a|^p + |b|^p \geq 2^{1-p}|a+b|^p$ we obtain

$$\begin{aligned} I_1 + I_2 &\geq \frac{1}{2} \int_{1/2}^1 \int_{B_t} \left(|u(x) - u_{B_t}|^p + \frac{2^{-p}|B_{1/2}|}{|B_1|} |u_{B_t}|^p \right) dx \nu(dt) \\ &\geq \frac{2^{-p}|B_{1/2}|}{2|B_1|} 2^{1-p} \int_{1/2}^1 \int_{B_t} |u(x)|^p dx \nu(dt) \\ &= \frac{|B_{1/2}|}{|B_1|} 2^{-2p} \int_{B_1} |u(x)|^p \phi(x) dx \end{aligned}$$

and the proof is finished. \square

3. APPLICATIONS

Let us discuss some corollaries. Corollary 3 is well-known [5], although our class of admissible weights is more general. Proposition 4 allows to deduce a weighted Poincaré inequality for fractional Sobolev norms from an unweighted version. Corollaries 5 and 6 give a more concrete result for fractional Sobolev norms. The first allows for more general kernels and exponents p . Corollary 6 improves [2, Theorem 5.1] because the result is robust for $s \rightarrow 1-$ and allows for general weights and exponents p .

Corollary 3. *Let $p \geq 1$ and ϕ be an admissible weight. Consider $X = \mathbb{R}^d$ equipped with the Lebesgue measure and the Euclidean metric. There exists a positive constant C depending on p, d and ϕ such that*

$$(7) \quad \int_{B_1} |u(x) - u_{B_1}^\phi|^p \phi(x) dx \leq C \int_{B_1} |\nabla u(x)|^p \phi(x) dx,$$

for every $u \in W^{1,p}(B_1)$.

Proposition 4. *Let $p \geq 1$ and let ϕ be an admissible weight of the form $\phi = \Phi(\rho(\cdot, x_0))$. Assume that for some kernel $k : B_1 \times B_1 \rightarrow [0, \infty)$ and some positive constant C the following inequality holds*

$$(8) \quad \int_{B_r} |u(x) - u_{B_r}|^p dx \leq C \int_{B_r} \int_{B_r} |u(x) - u(y)|^p k(x, y) dy dx,$$

whenever $r \in (\frac{1}{2}, 1]$ and $u \in L^p(X)$. Then with $M = \frac{8^p |B_1|}{|B_{1/2}|} \frac{\Phi(0)}{\Phi(1/2)}$

$$(9) \quad \int_{B_1} |u(x) - u_{B_1}^\phi|^p \phi(x) dx \leq CM \int_{B_1} \int_{B_1} |u(x) - u(y)|^p k(x, y) (\phi(y) \wedge \phi(x)) dy dx$$

for $u \in L^p(X)$.

Corollary 5. *Let ϕ be an admissible weight of the form $\phi = \Phi(\rho(\cdot, x_0))$ and $p \geq 1$. Let $k : B_1 \times B_1 \rightarrow [0, \infty)$ be a kernel satisfying $k \geq c$ for some constant $c > 0$. There is a positive constant M depending on d, p and Φ such that for $u \in L^p(X)$*

$$(10) \quad \int_{B_1} |u(x) - u_{B_1}^\phi|^p \phi(x) dx \leq \frac{M}{c} \int_{B_1} \int_{B_1} |u(x) - u(y)|^p k(x, y) (\phi(y) \wedge \phi(x)) dy dx$$

for $u \in L^p(X)$.

Corollary 6. *Let $p \geq 1$, $r \geq 1$ and $0 < s_0 \leq s < 1$. Consider $X = \mathbb{R}^d$ equipped with the Lebesgue measure and the Euclidean metric. Let ϕ be an admissible weight of the form $\phi = \Phi(|\cdot|)$. Then there exists a positive constant C depending on p, d, s_0 and Φ such that*

$$(11) \quad \int_{B_1} |u(x) - u_{B_1}^\phi|^p \phi(x) dx \leq C(1-s) r^{p(1-s)} \int_{B_1} \int_{B_1} \frac{|u(x) - u(y)|^p}{|x - y|^{d+ps}} \chi_{\{|x-y| \leq \frac{1}{r}\}} (\phi(y) \wedge \phi(x)) dy dx$$

for all $u \in L^p(B_1)$.

Proof of Corollary 3. It is well-known that the following Poincaré inequality holds

$$(12) \quad \int_{B_r} |u(x) - u_{B_r}|^p dx \leq c r^p \int_{B_r} |\nabla u(x)|^p dx$$

for every $u \in W^{1,p}(B_r)$ and $r > 0$ where $c > 0$ depends on p and d . Set

$$F(u, r) = c r^p \int_{B_r} |\nabla u(x)|^p dx,$$

for $u \in W^{1,p}(B_1)$ and $F(u, r) = \infty$ otherwise. Then for $u \in W^{1,p}(B_1)$

$$\begin{aligned} \int_{1/2}^1 F(u, t) \nu(dt) &= c \int_{1/2}^1 t^p \int_{B_1} |\nabla u(x)|^p \chi_{B_t}(x) dx \nu(dt) \\ &\leq c \int_{B_1} |\nabla u(x)|^p \int_{1/2}^1 \chi_{B_t}(x) \nu(dt) dx = c \int_{B_1} |\nabla u(x)|^p \phi(x) dx. \end{aligned}$$

By Theorem 1 the assertion follows with $C = 2^{3p+d} \frac{\Phi(0)}{\Phi(1/2)} c$. \square

Proof of Proposition 4. Let

$$F(u, r) = C \int_{B_r} \int_{B_r} |u(x) - u(y)|^p k(x, y) dy dx.$$

Then

$$\begin{aligned} \int_{1/2}^1 F(u, t) \nu(dt) &= C \int_{1/2}^1 \int_{B_1} \int_{B_1} |u(x) - u(y)|^p k(x, y) \chi_{B_t}(y) \chi_{B_t}(x) dy dx \nu(dt) \\ &= C \int_{B_1} \int_{B_1} |u(x) - u(y)|^p k(x, y) \int_{1/2}^1 \chi_{B_t}(y) \chi_{B_t}(x) \nu(dt) dy dx \\ &= C \int_{B_1} \int_{B_1} |u(x) - u(y)|^p k(x, y) (\phi(y) \wedge \phi(x)) dy dx. \end{aligned}$$

The assertion now follows from Theorem 1. \square

Lemma 7. *Let $r \geq 1$, $p \geq 1$ and $0 < s < 1$. Then*

$$(13) \quad \int_{B_1} \int_{B_1} \frac{|u(x) - u(y)|^p}{|x - y|^{d+ps}} dy dx \leq (3r)^{p(1-s)} \int_{B_1} \int_{B_1} \frac{|u(x) - u(y)|^p}{|x - y|^{d+ps}} \chi_{\{|x-y| \leq \frac{1}{r}\}} dy dx$$

for all $u \in L^p(B_1)$.

Proof. Let n be a natural number such that $n \geq 2r > n - 1$. We introduce

$$A_k = A_k(x, y) = \frac{k}{n}y + \frac{n-k}{n}x, \quad k = 0, 1, \dots, n.$$

Then

$$\begin{aligned} I &= \int_{B_1} \int_{B_1} \frac{|u(x) - u(y)|^p}{|x - y|^{d+ps}} dy dx = \int_{B_1} \int_{B_1} \frac{|\sum_{k=1}^n (u(A_{k-1}) - u(A_k))|^p}{|x - y|^{d+ps}} dy dx \\ &\leq n^{p-1} \sum_{k=1}^n \int_{B_1} \int_{B_1} \frac{|u(A_{k-1}) - u(A_k)|^p}{|x - y|^{d+ps}} dy dx. \end{aligned}$$

Note that $|A_{k-1} - A_k| = \frac{1}{n}|x - y|$. If we substitute $\tilde{x} = A_{k-1}$, $\tilde{y} = A_k$, then $d\tilde{y} d\tilde{x} = n^{-d} dy dx$ (which follows by an elementary calculation, see also [3, page 570]). Moreover, $\tilde{x}, \tilde{y} \in B_1$

with $|\tilde{x} - \tilde{y}| \leq \frac{2}{n} \leq \frac{1}{r}$. Hence

$$I \leq n^{p-ps} \int_{B_1} \int_{B_1} \frac{|u(\tilde{x}) - u(\tilde{y})|^p}{|\tilde{x} - \tilde{y}|^{d+ps}} \chi_{\{|\tilde{x}-\tilde{y}| \leq \frac{1}{r}\}} d\tilde{y} d\tilde{x}.$$

Since $n < 2r + 1 \leq 3r$, the assertion follows. \square

Proof of Corollary 5. First we use a well-known argument to obtain a nonweighted Poincaré inequality. By the convexity of the function $x \mapsto |x|^p$ it holds $|a+b|^p \geq |a|^p + bp|a|^{p-1} \operatorname{sgn}(a)b$. Thus

$$\begin{aligned} \int_{B_r} \int_{B_r} |u(x) - u(y)|^p k(x, y) dy dx &\geq c \int_{B_r} \int_{B_r} |(u(x) - u_{B_r}) + (u_{B_r} - u(y))|^p dy dx \\ &\geq c|B_r| \int_{B_r} |u(x) - u_{B_r}|^p dx \\ &\geq c|B_{1/2}| \int_{B_r} |u(x) - u_{B_r}|^p dx, \end{aligned}$$

whenever $u \in L^p(B_r)$ and $\frac{1}{2} < r \leq 1$.

The assertion follows now from Proposition 4. \square

Proof of Corollary 6. From [4] and [1, page 80] we know that there exists a constant $C = C(p, d, s_0)$, such that for $s_0 \leq s < 1$

$$(14) \quad \int_{B_r} |u(x) - u_{B_r}|^p dx \leq C(1-s)r^{ps} \int_{B_r} \int_{B_r} \frac{|u(x) - u(y)|^p}{|x-y|^{d+ps}} dy dx,$$

for all $u \in L^p(B_1)$. The assertion follows now from (14), Proposition 4 and Lemma 7. \square

REFERENCES

- [1] J. Bourgain, H. Brezis, and P. Mironescu. Limiting embedding theorems for $W^{s,p}$ when $s \uparrow 1$ and applications. *J. Anal. Math.*, 87:77–101, 2002. Dedicated to the memory of Thomas H. Wolff.
- [2] Z.-Q. Chen, P. Kim, and T. Kumagai. Weighted Poincaré inequality and heat kernel estimates for finite range jump processes. *Math. Ann.*, 342(4):833–883, 2008.
- [3] B. Dyda. On comparability of integral forms. *J. Math. Anal. Appl.*, 318(2):564–577, 2006.
- [4] A. C. Ponce. An estimate in the spirit of Poincaré’s inequality. *J. Eur. Math. Soc. (JEMS)*, 6(1):1–15, 2004.
- [5] L. Saloff-Coste. *Aspects of Sobolev-type inequalities*, volume 289 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2002.

B.D. AND M.K.: FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT BIELEFELD, POSTFACH 100131, D-33501 BIELEFELD,

B.D.: INSTITUTE OF MATHEMATICS AND COMPUTER SCIENCE, WROCLAW UNIVERSITY OF TECHNOLOGY, WYBRZEŻE WYSPIAŃSKIEGO 27, 50-370 WROCLAW, POLAND

E-mail address: bdyda (at) pwr wroc pl dyda (at) math uni-bielefeld de

E-mail address: moritz.kassmann (at) uni-bielefeld de

URL: www.math.uni-bielefeld.de/~kassmann